

A Guide to the  
**AMPHIBIANS & REPTILES**  
of SOPB Group of Tinbarap Estates  
and Adjacent Regions, Sarawak



Indraneil Das, Ivan Chiron Yaman, Kho Lip Khoon  
Pui Yong Min, Adi Shabranji, Izneil Nashriq and Anthony Pine





*Jointly published by*

**Institute of Biodiversity and Environmental Conservation  
Universiti Malaysia Sarawak  
94300 Kota Samarahan Sarawak, Malaysia**



**Malaysian Palm Oil Board  
No 6, Persiaran Institusi, Bandar Baru Bangi  
43000 Kajang, Selangor Darul Ehsan, Malaysia**



**Sarawak Oil Palms Berhad  
No. 124-126, Jalan Bendahara  
P.O. Box 547, 98007 Miri, Sarawak**

Copyright © 2019 Institute of Biodiversity and Environmental Conservation  
Universiti Malaysia Sarawak.


All rights reserved. No part of this publication may be reproduced, stored in a retrieval form or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission of the Publisher.

The views expressed are those of the respective authors, and not necessarily of those of the Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak.

*A Guide to the Amphibians and Reptiles of SOPB Group Tinbarap Estates and Adjacent Regions, Sarawak* by Indraneil Das, Ivan Chiron Yaman, Kho Lip Khoon, Pui Yong Min, Adi Shabranji, Izneil Nashriq and Anthony Pine.

ISBN 978-967-5418-72-3

Printed by Lee Ming Press Sdn Bhd, Kuching, Sarawak, Malaysia.



#### **Front Cover**

*Cuora amboinensis* (top)

*Trimeresurus borneensis* (bottom, left)

*Pulchrana baramica* (bottom, right)

Photos by Indraneil Das

#### **Back Cover**

Map showing sonar area mapped and crocodile sightings at Sungai Bakong, including zoomed views of sonar mosaics and depth recordings  
Photo by Anthony Pine

#### **Concept and Design**

Pui Yong Min & Indraneil Das

# **A Guide to the Amphibians and Reptiles of SOPB Group of Tinbarap Estates and Adjacent Regions, Sarawak**

*by*

Indraneil Das<sup>1</sup>, Ivan Chiron Yaman<sup>2</sup>, Kho Lip Khoon<sup>2</sup>  
Pui Yong Min<sup>1</sup>, Adi Shabranji<sup>1</sup>, Izneil Nashriq<sup>1</sup> and Anthony Pine<sup>1</sup>

<sup>1</sup>Institute of Biodiversity and Environmental Conservation  
Universiti Malaysia Sarawak

<sup>2</sup>Biodiversity and Conservation Unit  
Sarawak Oil Palms Berhad  
(Tinbarap Group of Estates)



**2019**



*Polypedates colletti* (Collett's Tree Frog)

# INTRODUCTION

Tinbarap Group of Estates, under the Sarawak Oil Palms Berhad, owns a total of 42 estates, spread along the lowlands of northern and central Sarawak, in East Malaysia.

Among its green achievements is the creation and maintenance of a High Conservation Value (HCV) forest patch, a reservoir of local biodiversity and an area where intensive research and monitoring of plants and animals can take place.

This guide covers the amphibians and reptiles of the site, based on rapid assessments carried out between 2016-2018. The Tinbarap HVC forest and adjacent areas are home to six species of amphibians and 12 species of reptiles, several of which are endemic to Borneo.

This field guide to the amphibians and reptiles of region, and is based on our inventories. The cut-off date for the checklist was 5 September 2019. A total of 18 species are described in this work, representing a generalised lowland, especially, peat swamp, herpetofauna of Borneo. For all species, we provide a common (English) name, the current scientific name and authority, size information (maximum snout-vent length for frogs, lizards and snakes and carapace length for turtle), brief description of external morphology and colouration of the adult stages relevant to its identification. For each species, we present an image of an adult.

Further information can be obtained from the references listed at the end of this field guide.

Also appended to this guide are recommended actions for snake bites and suggestions to reduce human-crocodile conflict at the site.

# BUFONIDAE

*Ingerophrynus quadriporcatus* (Boulenger, 1887)    Size 60 mm  
Four-ridged Toad

## Identification

Body stocky; a pair of ridges between eyes; parotoid gland over three times as long as wide, followed by rows of warts; fingers not fully webbed; back orange or yellow-brown; belly cream coloured.



# DICROGLOSSIDAE

*Fejervarya limnocharis* (Gravenhorst, 1829)  
Cricket Frog

Size 60 mm

## Identification

Body moderately slender; snout narrow; back with broken longitudinal ridges; outer metatarsal tubercle absent; toes with pointed tips, less than half webbed; back brown to brownish-grey, some with a pale yellow vertebral line; males with a black, M-shaped band across throat; belly cream-coloured.



# DICROGLOSSIDAE

*Limnonectes paramacrodon* (Inger, 1966)  
Masked Frog

Size 75 mm

## Identification

Body robust; snout rounded; tympanum and supratympanic fold distinct; Back mostly smooth, with a few low tubercles; toe tips widened and webbed to tips, except toe IV, with one phalange free; males lack vocal sacs; back greyish-brown; dark mask covers tympanum and supratympanic fold; belly yellow.



# RANIDAE

*Pulchrana baramica* (Boettger, 1900)  
Brown Marsh Frog

Size 71 mm

## Identification

Body slender; skin granular to slightly warty; head relatively large; tympanum visible; toes about half webbed; fingers long; upper lip with pale spots; back mid-brown to dark brown, with darker spots on flanks; tympanum dark brown, with a distinct pale centre; belly cream or pale brown.



# RHACOPHORIDAE

*Polypedates colletti* (Boulenger, 1890)\*  
Collett's Tree Frog

Size 78 mm

## Identification

Body slender; head triangular; snout pointed; tympanum distinct; most toes extensively webbed; back and belly coarsely granular; a conical appendage on heel; back tan to reddish-brown or brownish-grey; a dark hour-glass shaped pattern; flanks with scattered dark spots; limbs with dark bars.



Hour-glass shaped pattern

# RHACOPHORIDAE

*Polypedates otilophus* (Boulenger, 1893)\*  
File-eared Tree Frog

Size 97 mm

## Identification

Body robust; a bony ridge over tympanum; head large, triangular; tips of fingers and toes disk-like; toes half webbed; a fleshy projection at heel; back and forehead bright yellow to yellowish brown, with thin dark lines; belly cream-coloured.



# CROCODYLIDAE

*Crocodylus porosus* Schneider, 1801

**Size** 6.7 m

Saltwater Crocodile

## Identification

Head large with a heavy snout; a pair of ridges running from orbit to the centre of snout; scales on back more oval than in most other crocodiles; anterior nuchal scales usually absent; juveniles brightly coloured, being black spotted or blotched on a pale yellow or grey background; adults greyish-olive, bands showing less contrast



# GEOEMYDIDAE



*Cuora amboinensis* (Riche in: Daudin, 1801)  
Malayan Box Turtle

**Size** 216 mm

## **Identification**

Shell high-domed and smooth, with a single keel in adults, juveniles with two additional keels; carapace olive, brown or nearly black; plastron yellow or cream, with a single black blotch; face yellow-striped.

# AGAMIDAE

*Bronchocela cristatella* (Kuhl, 1820)

Crested Green Lizard

**Size** 130 mm

## Identification

Body slender, compressed; head long; nuchal crest short, continuous with dorsal crest; gular sac small; tympanum large; dorsal scales smooth; back green, sometimes with white or light blue spots that may form bars, or with white bars, changeable to brown; pale grey stripe across eyes; belly yellowish-green.



## GEKKONIDAE



*Cyrtodactylus pubisulcus* Inger, 1958\*  
Grooved Bent-toed Gecko

Size 77 mm

### Identification

Body slender; skin fold on flanks present; precloacal groove present; dorsal tubercles in irregular longitudinal rows; dorsum grey or brown, with dark cross-bars or blotches, sometimes arranged in a longitudinal series; belly cream coloured.

# GEKKONIDAE



*Gekko smithii* Gray, 1842  
Smith's Giant Gecko

**Size** 191 mm

## Identification

Body robust; head large; dorsum with scattered tubercles; precloacal pores in short angular series; dorsum greyish-brown, with series of white spots forming bands, tail dark-banded; belly cream coloured, with grey patches; eye green.

## GEKKONIDAE



*Hemidactylus frenatus* Duméril & Bibron, 1836  
Asian House Gecko

Size 67 mm

### Identification

Body robust, slightly flattened; head large; tail segmented; dorsal scales smooth; sides of tail with enlarged tubercles; dorsum greyish-brown, with darker markings; light brown streak across sides of head; belly cream coloured.

## SCINCIDAE

*Eutropis multifasciatus* (Kuhl, 1820)  
Common Ground Skink

**Size** 137 mm

### Identification

Body robust; head distinct; lower eyelid scaly; paired enlarged nuchals; auricular opening small, with small, pointed lobules; dorsal scales with three (rarely five) keels; dorsum brown with a yellow or red stripe or a series of white spots or streaks along flanks; belly cream coloured.



## SCINCIDAE

*Sphenomorphus cyanolaemus* Inger & Hosmer, 1965\* **Size** 60 mm  
Blue-bellied Litter Skink

### Identification

Body slender; limbs and tail long; snout obtuse; ear opening lacking lobules; scale smooth; head, throat and chest of males deep blue, in females, light blue; and lower abdomen pale blue in males, yellowish-orange in female; back bronze-brown to olive-brown, with a series of yellowish-brown blotches along flanks.



# **VARANIDAE**

*Varanus salvator* (Laurenti, 1768)  
Water Monitor Lizard

**Size** 910 mm

## **Identification**

Body robust; snout depressed; nostril rounded or oval; tail strongly compressed with a double-toothed crest above; juveniles dark dorsally, yellow spotted or ocelli in transverse series; snout black barred; belly yellow with narrow, and black vertical V-shaped marks.



# CROTALIDAE

*Trimeresurus borneensis* (Peters, 1872)\*  
Bornean Flat-nosed Pit Viper

Size 830 mm

## Identification

Back mottled brown, with a dark brown saddle-like pattern comprising 20–30 blotches or cross-bars, or even bright yellow with darker mottling; a pale stripe present from eye to neck.



## CROTALIDAE

*Tropidolaemus subannulatus* (Gray, 1842)  
Bornean Keeled Green Pit Viper

Size 963 mm

### Identification

Body slender in juveniles, relatively thick in adults; head distinct from neck; tail prehensile; scales keeled; back green or greenish-blue, with dark cross-bars blue and white, red and white, blue, blue and red or white in adult females; white spots or white and red spots in males and juveniles; pale stripe from eyes to back of head in adult females, white and red stripe in juveniles and males; belly unpatterned.



# PYTHONIDAE

*Malayopython reticulatus* (Schneider, 1801)

**Size** > 10 m

Reticulated Python

## Identification

Body relatively elongated and slender; head distinct from neck; pupil vertical; anal entire; dorsum yellow or brown, with dark, rhomboidal markings; black median line from snout to nape; belly yellow with small brown spots.



# SNAKE-BITE MANAGEMENT

Two venomous snakes were recorded within the Estate, and we provide brief notes on these species, venom and suggestions for treatment for their bites. Additional venomous snakes are expected to occur, especially the Spitting Cobra, *Naja sumatrana*.

## *Tropidolaemus subannulatus*


### Bornean Keeled Green Pit Viper

**Local names:** Bahasa Malaysia *Ular Kapak*; Iban: *Ular Engkaradau*

**Where Found:** Usually in low vegetation and mid-level of trees.

**Venom and Treatment:** Venom is a mixture of procoagulants; anticoagulants; additionally, haemorrhagins and necrotoxins are possibly present. Envenoming may be lethal and patients need to be taken immediately to Miri Hospital for observation and if required, antivenom. The appropriate serum is the Green Pit Viper Antivenin (manufactured by the Thai Red Cross Society).





*Trimeresurus borneensis*

**Bornean Palm Pit Viper**

**Local Name:** Unrecorded.

**Where Found.** Young ones are found on the ground, while adults can climb bushes and low trees.

**Venom and Treatment:** Venom contains procoagulants and haemorrhagins, and necrotoxins are possibly present. Bites have the potential for serious envenoming and require immediate medical attention at the Miri General Hospital. The Green Pit Viper Antivenin (manufactured by the Thai Red Cross Society) is recommended for its bite, after examination of the patient by a qualified medical practitioner.



# AVOIDING CROCODILE CONFLICT

- \* From our surveys throughout drainages and water sources within the area, it is unlikely that crocodiles would pose a major threat to safety.
- \* Drainages within the Estate are relatively shallow and do not offer resources necessary to support a significant crocodile population. Nonetheless, small crocodiles do make move into plantation areas.
- \* For rare occasions when adult crocodiles enter plantation areas, it can pose significant threat to safety of plantation personnel.
- \* Workers should be encouraged to report crocodile sightings of subadults or adults to plantation managers. Plantation staff would then be recommended to contact a biologist or surveyor to confirm the sighting using night-time eye-shine surveys.
- \* Recommendations could then be made on removal of the animal from the plantation zone.
- \* Crocodile safety and awareness should be included in briefings to workers and personnel that operate closer to major water bodies, such as Sungai Karap and Sungai Bakong.
- \* The southern area of Tinbarap Estate, adjacent to Miri Timur plantation estates, does have a shared drainage area that has reports of crocodiles by security staff. This area likely has the highest chance of crocodiles entering into the estate area.
- \* It is recommended that workers be aware that crocodiles may exist and that to be careful during areas in close proximity to major drainage areas or during night-time (i.e., fishing activities). Any food should also be disposed of properly in this area, as to avoid attracting crocodiles to these drainages.
- \* Although crocodile appear to pose an insignificant threat to safety at Tinbarap Estate, it would be recommended to maintain records and reporting procedures for crocodile sightings if they occur.

# ACKNOWLEDGEMENTS

This is a collaborative project led by Malaysian Palm Oil Board (MPOB). We thank Sarawak Oil Palms Berhad for supporting the project through a research grant. The SOPB assisted with all aspects of project planning and field work. In particular, we would like to thank Mr. Paul Wong, GCEO, Mr. Chua Kian Hong, Group Plantation Controller, Mr. Jiw William, Tinbarap 4 Estate Manager, Mr. Junis Jampang Aton, Tinbarap 5 EM, and Mr. Apong Siri, Tinbarap 7 SEM.

The project was administered by the Research and Innovative Management Centre, Universiti Malaysia Sarawak, for which we thank Prof Dr Lo May Chiun and her staff. Lab and office space was provided by the Institute of Biodiversity and Environmental Conservation, UNIMAS, for which we thank our Director, Prof. Dr. Gabriel Tonga Noweg, and administrative staff, who helped with the printing of the field guide, including Ms. Felicia Anthony Reyap, Pasey Lisus and Ketty Daun.



*The habitat adjacent to the Sungai Karap bridge during daytime*

## **REFERENCES**

- Das, I. 2010.** A field guide to the reptiles of south-east Asia. New Holland Publishers (UK), Ltd., London. 376 pp.
- Frost, D.R. 2019.** Amphibian species of the world: an online reference. Version 6.0 (5 September 2019). Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.php>. American Museum of Natural History, New York, USA.
- Haas, A., I. Das, S. Hertwig, Y.M. Pui & A. Jankowski. 2019.** Frogs of Borneo. Frogs and tadpoles of East Malaysia. Inter-net resource at: <http://frogssofarne.org>
- Hazebroek, H.P. & A.K.A. Morshidi. 2000.** National Parks of Sarawak. Natural History Publications (Borneo) Sdn. Bhd, Kota Kinabalu. xii + 503 pp.
- Inger, R.F. 1966.** The systematics and zoogeography of the Amphibia of Borneo. *Fieldiana Zoology* 52:1–402. Reprinted 1990 and 2005, by Lun Hing Trading, Kota Kinabalu and Natural History Publications (Borneo) Sdn. Bhd., Kota Kinabalu, respectively.
- Inger, R.F., R.B. Stuebing, T.U. Gafe & J.M. Dehling. 2017.** A field guide to the frogs of Borneo. Third Edition. Natural History Publications (Borneo) Sdn Bhd., Kota Kinabalu. viii + 228 pp.



▲ *Trimeresurus borneensis*



